

Molecular Basis of Cholera Toxin Action in Mammalian Cells: Modulation of Toxin ADP-ribosyltransferase Activity by Endogenous ADP-ribosylation Cycles

Joel Moss, Jianfeng Zhu, Shunya Oka, and Jiro Kato

Cholera toxin, produced by *Vibrio cholerae*, is responsible, in large part, for the devastating diarrhea characteristic of cholera. The toxin exerts its effects on cells through the ADP-ribosylation of Gas, the stimulatory guanine nucleotide-binding protein that activates adenylyl cyclase, resulting in increased intracellular cyclic AMP content, and the often life-threatening derangements in fluid and electrolyte fluxes that are characteristic of the disease (1). Mammalian cells contain NAD:arginine ADP-ribosyltransferases, which, similar to cholera toxin, catalyze the modification of arginine residues in proteins; the ADP-ribose-arginine bonds are cleaved by ADP-ribose-(arginine)protein hydrolases (ARH), which regenerate the free (arginine)protein, thus completing an ADP-ribosylation cycle (2). We identified three genes that encode proteins with ARH-like sequences (3). Only one of the protein products of these genes, ARH1, was capable of cleaving ADP-ribose-(arginine)protein (3). We hypothesized that ARH1, if capable of cleaving ADP-ribose(arginine)Gas, synthesized by cholera toxin, might reverse or limit the effects of cholera toxin on intestinal cells. In ARH1, aspartate residues 60 and 61 are critical for activity; their replacement with alanine results in an ARH1 protein with <0.1% of wild-type activity (4). ARH1 knockout mice were generated by replacing the exon containing the critical aspartate residues with a neomycin cassette. The ARH1 knockout mice are viable. Effects of cholera toxin on fluid accumulation in intestinal loops were significantly greater in the knockout than in the wild-type animals. Effects of cyclic AMP, the downstream signaling molecule generated by cholera toxin-catalyzed ADP-ribosylation were not, however, different in intestinal loops of the wild-type and knockout animals. Cholera toxin-generated ADP-ribosylarginine content and Gas modification were significantly higher in intestinal epithelial cells from the knockout than from the wild-type mice. Similar findings were obtained with knockout and wild-type cells in culture. Thus, the effects of cholera toxin were significantly influenced by cellular ARH1 activity, which could function as a modifier gene in disease and serve as a potential therapeutic target. Presumably, in patients with cholera, toxin activity overwhelms this novel host defense, leading to persistent activation of adenylyl cyclase and the diarrheal syndrome characteristic of the disease.

(1) Moss J, Vaughan M. ADP-ribosylation of guanyl nucleotide-binding regulatory proteins by bacterial toxins. *Adv Enzymol Relat Areas Mol Biol.* 1988;61:303-79.

(2) Okazaki IJ, Moss J. Glycosylphosphatidylinositol-anchored and secretory isoforms of mono-ADP-ribosyltransferases. *J Biol Chem.* 1998;273(37):23617-20.

(3) Oka S, Kato J, Moss J. Identification and characterization of a mammalian 39-kDa poly(ADP-ribose) glycohydrolase. *J Biol Chem.* 2006;281(2):705-13.

(4) Konczalik P, Moss J. Identification of critical, conserved vicinal aspartate residues in mammalian and bacterial ADP-ribosylarginine hydrolases. J Biol Chem. 1999;274(24):16736-40.